

Parametric Resonant Phenomena in Bose-Einstein Condensates: Breaking of Macroscopic Quantum Self-Trapping

Luca Salasnich

Istituto Nazionale per la Fisica della Materia, Unità di Milano,

Dipartimento di Fisica, Università di Milano,

Via Celoria 16, 20133 Milano, Italy

E-mail: salasnich@mi.infm.it

Abstract

We analyze the periodic tunneling of a Bose-Einstein condensate in a double-well potential which has an oscillating energy barrier. We show that the dynamics of the Bose condensate critically depends on the frequency ω of the oscillating energy barrier. In the regime of periodic macroscopic quantum tunneling (PMQT) with frequency ω_J , the population imbalance of the condensate in the two wells can be enhanced under the condition of parametric resonance $\omega = 2\omega_J$. Instead, in the regime of macroscopic quantum self-trapping (MQST), we find that MQST can be reduced or suppressed under the condition of parametric resonance between the frequency ω of the energy barrier and the frequency ω_{ST} of oscillation through the barrier of the very small fraction of particles which remain untrapped during MQST.

I. INTRODUCTION

Macroscopic quantum tunneling with dilute Bose-Einstein condensates of alkali-metal atoms has been the subject of many theoretical [1-4] and experimental [5-7] studies. Here we consider the problem of trapped Bosons in a double-well potential and investigate the effect of a periodically varying barrier in the tunneling of the Bose condensate. We study the problem using the two-mode classical-like equations [1,2] and find that remarkable effects are related to parametric resonance [8] both in the periodic macroscopic quantum tunneling regime and in the macroscopic quantum self-trapping regime.

II. TUNNELING IN A DOUBLE-WELL POTENTIAL

In a dilute gas of N Bosons at zero temperature practically all particles are in the same single-particle state of the density matrix $\rho(\mathbf{r}, \mathbf{r}'; t)$ [9]. This macroscopically occupied single-particle state is called Bose-Einstein condensate and its wavefunction $\psi(\mathbf{r}, t)$ is well described by the Gross-Pitaevskii equation

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}, t) = \left[-\frac{\hbar^2}{2m} \nabla^2 + U(\mathbf{r}) + g|\psi(\mathbf{r}, t)|^2 \right] \psi(\mathbf{r}, t) , \quad (1)$$

where $U(\mathbf{r})$ is the external potential, $g = 4\pi\hbar^2 a_s/m$ is the inter-atomic strength with a_s the s-wave scattering length, and the wavefunction is normalized to N [10].

Let us consider a double-well external potential $U(\mathbf{r})$. In the tunneling regime, the wavefunction of the Bose-condensate can be approximated in the following way

$$\psi(\mathbf{r}, t) = \sqrt{N_1(t)} e^{i\phi_1(t)} \psi_1(\mathbf{r}) + \sqrt{N_2(t)} e^{i\phi_2(t)} \psi_2(\mathbf{r}) , \quad (2)$$

where $N_1(t)$ is the number of particles in the first well and $N_2(t)$ is the number of particles in the second well, such as $N = N_1(t) + N_2(t)$ [1]. If the double-well is symmetric, it is not difficult to show that the time-dependent behavior of the condensate in the tunneling energy range can be described, with a suitable rescaling of time, by the two-mode equations

$$\dot{\zeta} = -\sqrt{1 - \zeta^2} \sin \phi , \quad \dot{\phi} = \Lambda \zeta + \frac{\zeta}{\sqrt{1 - \zeta^2}} \cos \phi , \quad (3)$$

where $\zeta = (N_1 - N_2)/N$ is the fractional population imbalance of the condensate in the two wells, $\phi = \phi_1 - \phi_2$ is the relative phase (which can be initially zero), and $\Lambda = 2E_I/E_T$ with E_I the inter-atomic energy and E_T the tunneling energy, i.e. the kinetic+potential energy splitting between the ground state and the quasi-degenerate odd first excited state of the GP equation [1,2]. Note that $E_T = \hbar\omega_0$, where ω_0 is the oscillation frequency of the Bose condensate between the two wells when the inter-atomic interaction is zero ($E_I = 0$) [1-4].

The two-mode equations have been studied by many authors and they are the Hamilton equations of the following Hamiltonian

$$H = \frac{\Lambda}{2}\zeta^2 - \sqrt{1 - \zeta^2} \cos \phi, \quad (4)$$

where ζ is the conjugate momentum of the generalized coordinate ϕ . Actually, it is possible to introduce [1] an equivalent Hamiltonian, given by

$$H' = \frac{1}{2}p_\zeta^2 + a\zeta^2 + b\zeta^4, \quad (5)$$

where

$$a = \frac{1}{2}\sqrt{1 - \Lambda H_0}, \quad b = \frac{\Lambda^2}{8}, \quad (6)$$

with $H_0 = \Lambda\zeta(0)^2/2 - \sqrt{1 - \zeta(0)^2} \cos \phi(0)$. For the Hamiltonian H' the generalized coordinate is ζ and its conjugate momentum is p_ζ . The equations of motion derived by H' are

$$\dot{\zeta} = p_\zeta, \quad \dot{p}_\zeta = -2a\zeta - 4b\zeta^3. \quad (7)$$

The analysis of the two-mode equations (3) or (7) has shown that for $\Lambda < 2$ there is a periodic macroscopic quantum tunneling (PMQT) of the Bose-condensate with Josephson-like oscillations of frequency $\omega_J = \omega_0\sqrt{1 + \Lambda}$ [1,2]. Instead, for $\Lambda > 2$, there exists a critical $\zeta_c = 2\sqrt{\Lambda - 1}/\Lambda$ such that for $0 < \zeta(0) \ll \zeta_c$ there is PMQT of condensate with frequency ω_J , but for $\zeta_c < \zeta(0) \leq 1$ there is macroscopic quantum self-trapping (MQST) of the condensate: even if the populations of the two wells are initially set in an asymmetric state ($\zeta(0) \neq 0$) they maintain, on the average, the original population imbalance with a very small periodic transfer of particles through the barrier with frequency $\omega_{ST} = \omega_0\sqrt{2(\Lambda H_0 - 1)}$ [1,2].

III. AN OSCILLATING BARRIER IN THE DOUBLE-WELL POTENTIAL

In this section we analyze the effect of a periodic oscillating energy barrier in the double-well potential. The presence of a oscillating barrier can be modelled by a time-dependent Λ . We choose the following form

$$\Lambda(t) = \Lambda_0(1 + \epsilon \sin(\omega t)) , \quad (8)$$

where Λ_0 is the static value, ϵ is a small perturbation ($\epsilon \ll 1$) and ω is the oscillation frequency of the barrier.

We numerically solve the two-mode equations (7) with (8) by using a forth-order Runge-Kutta algorithm. First we consider the case of pulsed macroscopic quantum tunneling (PMQT). We set $\Lambda_0 = 0.8$, $\zeta(0) = 0.5$ and $\phi(0) = 0$. In Figure 1 we plot the population imbalance (t, ζ) and its phase-space portrait (ζ, p_ζ) for some values of ϵ and ω . For $\epsilon = 0$ the population imbalance harmonically oscillates between the values -0.5 and 0.5 with frequency $\omega_J = 1.27$, but for $\epsilon \neq 0$ the motion is not fully harmonic. In particular, Figure 1 shows that under the parametric resonance condition $\omega = 2\omega_J$ the dynamics of the population imbalance is modified: the amplitude of the oscillation is modulated and reaches the values -1 and 1 .

Now we consider the case of macroscopic quantum self-trapping (MQST). We set $\Lambda_0 = 25$, $\zeta(0) = 0.6$ and $\phi(0) = 0$. Figure 2 confirms that for $\epsilon = 0$ the population imbalance does not change sign and it periodically oscillates between the values 0.47 and 0.6 with frequency $\omega_{ST} = 13.53$. As shown in Figure 2, MQST is strongly affected by the parametric resonance between the frequency ω_{ST} of MQST oscillation and the frequency ω of oscillation of the energy barrier. Moreover, Figure 3 shows that at the parametric resonance condition $\omega = 2\omega_{ST}$ and with a sufficiently large perturbation ($\epsilon = 0.2$) the system eventually escapes from the self-trapping configuration. Note that this remarkable effect has been confirmed [11] by using the nonpolynomial Schrödinger equation, an effective one-dimensional equation derived from the three-dimensional Gross-Pitaevskii equation [12].

CONCLUSIONS

We have studied the periodic tunneling and the quantum self-trapping of a Bose-Einstein condensate in a double-well potential with an oscillating energy barrier. We have used the two-mode classical-like equations to find that the periodic macroscopic quantum tunneling can be enhanced and the macroscopic quantum self-trapping can be reduced or suppressed by changing the frequency of the oscillating barrier. In the latter case, we have proved that the system escapes from the self-trapping configuration if the the frequency of oscillation of the double-well energy barrier and the frequency of MQST oscillations of the condensate satisfy the parametric resonance condition with a sufficiently large perturbation of the energy barrier.

REFERENCES

- [1] Smerzi, A., Fantoni, S., Giovannazzi, G. and Shenoy, S.R., 1997, Phys. Rev. Lett. **79** 4950; Raghavan, S., Smerzi, A., Fantoni, S., and Shenoy, S.R., 1999, Phys. Rev. A **59** 620.
- [2] Milburn, G.J., Corney, J., Wright, E., and Walls, D.F., 1997, Phys. Rev. A **55** 4318.
- [3] Zapata, I., Sols, F., and Leggett, A.J., 1998, Phys. Rev. A **57** R28.
- [4] Salasnich, L., Parola, A., and Reatto, L., 1999, Phys. Rev. A **60** 4171; Pozzi, B., Salasnich, L., Parola, A., and Reatto, L., 2000, Eur. Phys. J. D **11** 367.
- [5] Anderson, B.P. and Kasevich, M., 1999, Science **282** 1686.
- [6] Cataliotti, F.S., Burger, S., Fort, C., Maddaloni, P., Trombettoni, A., Smerzi, A., and M. Inguscio, 2001, Science **293** 843.
- [7] Morsch, O., Muller, J.H., Cristiani, M., Ciampini, D., and Arimondo, E., 2001, Phys. Rev. Lett. **87** 140402.

- [8] Landau, L.D. and Lifshits, E.M., 1991, *Mechanics, Course of Theoretical Physics*, vol. 3 (Pergamon Press, Oxford); Arnold, V.I., 1990, *Mathematical Methods of Classical Mechanics* (Springer, Berlin).
- [9] Courteille, P.W., Bagnato, V.S., and Yukalov, Y.I., 2001, *Laser Phys.* **11**, 659.
- [10] Gross, E.P., 1961, *Nuovo Cimento* **20** 454; Pitaevskii, L.P., 1961, *Zh. Eksp. Teor. Fiz.* **40**, 646.
- [11] Salasnich, L., Parola, A., and Reatto, L., 2002, *J. Phys. B* **35**, 3205.
- [12] Salasnich, L., 2002, *Laser Phys.* **12** 198; Salasnich, L., Parola, A., and Reatto, L., 2002, *Phys. Rev. A* **65** 043614.

FIGURES




FIG. 1. PMQT regime. Fractional population imbalance $\zeta(t)$ and its phase-space portrait (ζ, p_ζ) . Initial conditions: $\zeta(0) = 0.5$, $\phi(0) = 0$. $\Lambda = \Lambda_0(1 + \epsilon \sin(\omega t))$ where $\Lambda_0 = 0.8$. From top to bottom: (a) $\epsilon = 0$; (b) $\epsilon = 0.25$ and $\omega = \omega_J$; (c) $\epsilon = 0.25$ and $\omega = 2\omega_J$; where ω_J is the frequency of unperturbed PMQT oscillations.




FIG. 2. MQST regime. Fractional population imbalance $\zeta(t)$ and its phase-space portrait (ζ, p_ζ) . Initial conditions: $\zeta(0) = 0.6$, $\phi(0) = 0$. $\Lambda = \Lambda_0(1 + \epsilon \sin(\omega t))$ where $\Lambda_0 = 25$. From top to bottom: (a) $\epsilon = 0$; (b) $\epsilon = 0.1$ and $\omega = \omega_{ST}/2$; (c) $\epsilon = 0.1$ and $\omega = 2\omega_{ST}$; where ω_{ST} is the frequency of unperturbed MQST oscillations.




FIG. 3. MQST regime at the parametric resonance. Fractional population imbalance $\zeta(t)$ of the Bose condensate. Initial conditions: $\zeta(0) = 0.6$, $\phi(0) = 0$. $\Lambda = \Lambda_0(1 + \epsilon \sin(\omega t))$ where $\Lambda_0 = 25$ and $\omega = 2\omega_{ST}$ with ω_{ST} the frequency of unperturbed MQST oscillations. From top to bottom and from left to right: (a) $\epsilon = 0.01$; (b) $\epsilon = 0.05$; (c) $\epsilon = 0.1$; (d) $\epsilon = 0.2$.